


*Amendments to the Claims*

This listing of claims will replace all prior versions, and listings of claims in the application.

1. (Currently Amended) A composition comprising a synergistically effective ~~active compound~~ combination of compounds of the formula (I) (group 1)



in which

X represents halogen, alkyl, alkoxy, haloalkyl, haloalkoxy or cyano,

W¹, Y and Z independently of one another represent hydrogen, halogen, alkyl, alkoxy, haloalkyl, haloalkoxy or cyano,

A³ represents hydrogen, in each case optionally halogen-substituted alkyl, alkoxyalkyl, saturated, optionally substituted cycloalkyl in which optionally at least one ring atom is replaced by a heteroatom,


A⁴ represents hydrogen or alkyl, or

A³ and A⁴ furthermore together with the carbon atom to which they are attached represent a saturated or unsaturated, unsubstituted or substituted cycle which optionally contains at least one heteroatom,

D represents hydrogen or an optionally substituted radical selected from the group consisting of alkyl, alkenyl, alkoxyalkyl, and saturated cycloalkyl in which optionally one or more ring members are replaced by heteroatoms, or

A³ and D together with the atoms to which they are attached represent a saturated or unsaturated cycle which is unsubstituted or substituted in the A³,D moiety and optionally contains at least one heteroatom,


G¹ represents hydrogen (a) or represents one of the groups



- E represents a metal ion or an ammonium ion,  
 $\text{L}^1$  represents oxygen or sulfur,  
 $\text{M}^1$  represents oxygen or sulfur,  
 $\text{R}^{20}$  represents in each case optionally halogen-substituted alkyl, alkenyl, alkoxyalkyl, alkylthioalkyl, or polyalkoxyalkyl or optionally halogen-, alkyl- or alkoxy-substituted cycloalkyl which may be optionally interrupted by at least one heteroatom, or in each case optionally substituted phenyl, phenylalkyl, hetaryl, phenoxyalkyl or hetaryloxyalkyl,  
 $\text{R}^{21}$  represents in each case optionally halogen-substituted alkyl, alkenyl, alkoxyalkyl, or polyalkoxyalkyl or represents in each case optionally substituted cycloalkyl, phenyl or benzyl,  
 $\text{R}^{22}$  represents optionally halogen-substituted alkyl or optionally substituted phenyl,  
 $\text{R}^{23}$  and  $\text{R}^{24}$  independently of one another represent in each case optionally halogen-substituted alkyl, alkoxy, alkylamino, dialkylamino, alkylthio, alkenylthio, or cycloalkylthio or represent in each case optionally substituted phenyl, benzyl, phenoxy or phenylthio and  
 $\text{R}^{25}$  and  $\text{R}^{26}$  independently of one another represent hydrogen, in each case optionally halogen-substituted alkyl, cycloalkyl, alkenyl, alkoxy, or alkoxyalkyl, represent optionally substituted phenyl, represent optionally substituted benzyl or together with the nitrogen atom to which they are attached represent an optionally substituted ring which is optionally interrupted by oxygen or sulfur,


or an insecticidally active compound (group 2), selected from the group consisting of preferably

(2-1) amitraz (~~known from DE-A 20 61 132~~)




and/or

(2-2) buprofezin (~~known from DE-A 28 24 126~~)




and/or

(2-3) triazamate (~~known from EP-A 0 213 718~~)




and/or

(2-4) pymetrozine (~~known from EP-A 0 314 615~~)




and/or

(2-5) pyriproxyfen (~~known from EP-A 0 128 648~~)




and/or

(2-6) flonicamid (~~known from EP-A 0 580 374~~)




and/or and

(2-7) pirimicarb (known from GB 1 181 657)



and at least one active compound from the group of the anthranilamides  
anthranilamide of the formula (II)



in which

$A^1$  and  $A^2$  independently of one another represent oxygen or sulfur,

$X^1$  represents N or  $\text{CR}^{10}$ ,

$R^1$  represents hydrogen or represents  $\text{C}_1\text{-C}_6$ -alkyl,  $\text{C}_2\text{-C}_6$ -alkenyl,  $\text{C}_2\text{-C}_6$ -alkynyl or  $\text{C}_3\text{-C}_6$ -cycloalkyl, each of which is optionally mono- or polysubstituted, where the substituents independently of one another are selected from the group consisting of  $R^6$ , halogen, cyano, nitro, hydroxyl,  $\text{C}_1\text{-C}_4$ -alkoxy,  $\text{C}_1\text{-C}_4$ -alkylthio,  $\text{C}_1\text{-C}_4$ -alkylsulfinyl,  $\text{C}_1\text{-C}_4$ -alkylsulfonyl,  $\text{C}_2\text{-C}_4$ -alkoxycarbonyl,  $\text{C}_1\text{-C}_4$ -alkylamino,  $\text{C}_2\text{-C}_8$ -dialkylamino,  $\text{C}_3\text{-C}_6$ -cycloalkylamino, and  $(\text{C}_1\text{-C}_4\text{-alkyl})\text{-C}_3\text{-C}_6$ -cycloalkylamino and  $R^{11}$ ,

R<sup>2</sup> represents hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>2</sub>-C<sub>8</sub>-dialkylamino, C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino, C<sub>2</sub>-C<sub>6</sub>-alkoxycarbonyl or C<sub>2</sub>-C<sub>6</sub>-alkylcarbonyl,

R<sup>3</sup> represents hydrogen, R<sup>11</sup> or represents C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl, or C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, each of which is optionally mono- or polysubstituted, where the substituents independently of one another may be selected from the group consisting of R<sup>6</sup>, halogen, cyano, nitro, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>2</sub>-C<sub>6</sub>-alkoxycarbonyl, C<sub>2</sub>-C<sub>6</sub>-alkylcarbonyl, C<sub>3</sub>-C<sub>6</sub>-trialkylsilyl, R<sup>11</sup>, phenyl, phenoxy and a 5- or 6-membered heteroaromatic ring, where each phenyl, phenoxy and 5- or 6-membered heteroaromatic ring is optionally substituted and where the substituents independently of one another may be selected from one to three radicals W or one or more radicals R<sup>12</sup>, or

R<sup>2</sup> and R<sup>3</sup> may be attached to one another and form the ring M,

R<sup>4</sup> represents hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>2</sub>-C<sub>6</sub>-haloalkenyl, C<sub>2</sub>-C<sub>6</sub>-haloalkynyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl, halogen, cyano, nitro, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>2</sub>-C<sub>8</sub>-dialkylamino, C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino, or C<sub>3</sub>-C<sub>6</sub>-trialkylsilyl or represents phenyl, benzyl or phenoxy, each of which is optionally mono- or polysubstituted, where the substituents independently of one another are selected from the group consisting of C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>2</sub>-C<sub>4</sub>-alkenyl, C<sub>2</sub>-C<sub>4</sub>-alkynyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>2</sub>-C<sub>4</sub>-haloalkenyl, C<sub>2</sub>-C<sub>4</sub>-haloalkynyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl, halogen, cyano, nitro, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>2</sub>-C<sub>8</sub>-dialkylamino, C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino, C<sub>3</sub>-C<sub>6</sub>-(alkyl)cycloalkylamino, C<sub>2</sub>-C<sub>4</sub>-alkylcarbonyl, C<sub>2</sub>-C<sub>6</sub>-alkoxycarbonyl, C<sub>2</sub>-C<sub>6</sub>-alkylaminocarbonyl, C<sub>3</sub>-C<sub>8</sub>-dialkylaminocarbonyl and C<sub>3</sub>-C<sub>6</sub>-trialkylsilyl,

R<sup>5</sup> and R<sup>8</sup> in each case independently of one another represent hydrogen, or halogen or represent in each case optionally substituted C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, R<sup>12</sup>, G, J, -OJ, -OG, -S(O)<sub>p</sub>-J, -S(O)<sub>p</sub>-G, -S(O)<sub>p</sub>-phenyl, where the substituents independently of one another are selected from the group consisting of one to three radicals W, R<sup>12</sup>, C<sub>1</sub>-C<sub>10</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy and C<sub>1</sub>-C<sub>4</sub>-alkythio, where each substituent is optionally substituted by one or more substituents independently of one another selected from the group consisting of G, J, R<sup>6</sup>, halogen, cyano, nitro, amino, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>2</sub>-C<sub>8</sub>-dialkylamino, C<sub>3</sub>-C<sub>6</sub>-trialkylsilyl, phenyl and phenoxy, where each phenyl or phenoxy ring is optionally substituted and where the substituents independently of one another are selected from one to three radicals W or one or more radicals R<sup>12</sup>,

G in each case independently of one another represent a 5- or 6-membered non-aromatic carbocyclic or heterocyclic ring optionally contains one or two ring members from the group consisting of C(=O), SO and S(=O)<sub>2</sub> and which is optionally substituted by one to four substituents independently of one another selected from the group consisting of C<sub>1</sub>-C<sub>2</sub>-alkyl, halogen, cyano, nitro and C<sub>1</sub>-C<sub>2</sub>-alkoxy, or independently of one another represent C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl, C<sub>3</sub>-C<sub>7</sub>-cycloalkyl, (cyano)-C<sub>3</sub>-C<sub>7</sub>-cycloalkyl, (C<sub>1</sub>-C<sub>4</sub>-alkyl)-C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, or (C<sub>3</sub>-C<sub>6</sub>-cycloalkyl)-C<sub>1</sub>-C<sub>4</sub>-alkyl, where each cycloalkyl, (alkyl)cycloalkyl and (cycloalkyl)alkyl is optionally substituted by one or more halogen atoms,

J in each case independently of one another represent an optionally substituted 5- or 6-membered heteroaromatic ring, where the substituents independently of one another are selected from one to three radicals W or one or more radicals R<sup>12</sup>,

R<sup>6</sup> independently of one another represent -C(=E<sup>1</sup>)R<sup>19</sup>, -LC(=E<sup>1</sup>)R<sup>19</sup>, -C(=E<sup>1</sup>)LR<sup>19</sup>, -LC(=E<sup>1</sup>)LR<sup>19</sup>, -OP(=Q)(OR<sup>19</sup>)<sub>2</sub>, -SO<sub>2</sub>LR<sup>18</sup> or -LSO<sub>2</sub>LR<sup>19</sup>

where each E<sup>1</sup> independently of one another represents O, S, N-R<sup>15</sup>, N-OR<sup>15</sup>, N-N(R<sup>15</sup>)<sub>2</sub>, N-S=O, N-CN or N-NO<sub>2</sub>,

R<sup>7</sup> represents hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, halogen, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfonyl,

R<sup>9</sup> represents C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfinyl or halogen,

R<sup>10</sup> represents hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, halogen, cyano or C<sub>1</sub>-C<sub>4</sub>-haloalkoxy,

R<sup>11</sup> in each case independently of one another represents in each case optionally mono- to trisubstituted C<sub>1</sub>-C<sub>6</sub>-alkylthio, C<sub>1</sub>-C<sub>6</sub>-alkylsulfenyl, C<sub>1</sub>-C<sub>6</sub>-haloalkythio, C<sub>1</sub>-C<sub>6</sub>-haloalkylsulfenyl, phenylthio or phenylsulfenyl, where the substituents independently of one another are selected from the group consisting of W, -S(O)<sub>n</sub>N(R<sup>16</sup>)<sub>2</sub>, -C(=O)R<sup>13</sup>, -L(C=O)R<sup>14</sup>, -S(C=O)LR<sup>14</sup>, -C(=O)LR<sup>13</sup>, -S(O)<sub>n</sub>NR<sup>13</sup>C(=O)R<sup>13</sup>, -S(O)<sub>n</sub>NR<sup>13</sup>C(=O)LR<sup>14</sup> or and -S(O)<sub>n</sub>NR<sup>13</sup>S(O)<sub>2</sub>LR<sup>14</sup>,

L in each case independently of one another represents O, NR<sup>18</sup> or S,

R<sup>12</sup> in each case independently of one another represents -B(OR<sup>17</sup>)<sub>2</sub>, amino, SH, thiocyanato, C<sub>3</sub>-C<sub>8</sub>-trialkylsilyloxy, C<sub>1</sub>-C<sub>4</sub>-alkyl disulfide, -SF<sub>5</sub>, -C(=E<sup>1</sup>)R<sup>19</sup>, -LC(=E<sup>1</sup>)R<sup>19</sup>, -C(=E<sup>1</sup>)LR<sup>19</sup>, -LC(=E<sup>1</sup>)LR<sup>19</sup>, -OP(=Q)(OR<sup>19</sup>)<sub>2</sub>, -SO<sub>2</sub>LR<sup>19</sup> or -LSO<sub>2</sub>LR<sup>19</sup>,

Q represents O or S,

R<sup>13</sup> in each case independently of one another represent hydrogen or represent in each case optionally mono- or polysubstituted C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl or C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, where the substituents independently of one another are selected from the group consisting of R<sup>6</sup>, halogen, cyano, nitro, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>2</sub>-C<sub>8</sub>-dialkylamino, C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino and (C<sub>1</sub>-C<sub>4</sub>-alkyl)-C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino,

- R<sup>14</sup> in each case independently of one another represent in each case optionally mono- or polysubstituted C<sub>1</sub>-C<sub>20</sub>-alkyl, C<sub>2</sub>-C<sub>20</sub>-alkenyl, C<sub>2</sub>-C<sub>20</sub>-alkynyl or C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, where the substituents independently of one another are selected from the group consisting of R<sup>6</sup>, halogen, cyano, nitro, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>2</sub>-C<sub>8</sub>-dialkylamino, C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino and (C<sub>1</sub>-C<sub>4</sub>-alkyl)-C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino or represent optionally substituted phenyl, where the substituents independently of one another are selected from one to three radicals W or one or more radicals R<sup>12</sup>,
- R<sup>15</sup> in each case independently of one another represent hydrogen or represent in each case optionally mono- or polysubstituted C<sub>1</sub>-C<sub>6</sub>-haloalkyl or C<sub>1</sub>-C<sub>6</sub>-alkyl, where the substituents independently of one another are selected from the group consisting of cyano, nitro, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>2</sub>-C<sub>8</sub>-dialkylamino, C<sub>2</sub>-C<sub>6</sub>-alkoxycarbonyl, C<sub>2</sub>-C<sub>6</sub>-alkylcarbonyl, C<sub>3</sub>-C<sub>6</sub>-trialkylsilyl and optionally substituted phenyl, where the substituents independently of one another are selected from one to three radicals W or one or more radicals R<sup>12</sup>, or N(R<sup>15</sup>)<sub>2</sub> represents a cycle which forms the ring M,
- R<sup>16</sup> represents C<sub>1</sub>-C<sub>12</sub>-alkyl or C<sub>1</sub>-C<sub>12</sub>-haloalkyl, or N(R<sup>16</sup>)<sub>2</sub> represents a cycle which forms the ring M,
- R<sup>17</sup> in each case independently of one another represent hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, or B(OR<sup>17</sup>)<sub>2</sub>, represents a ring in which the two oxygen atoms are attached via a chain having two to three carbon atoms which are optionally substituted by one or two substituents independently of one another selected from the group consisting of methyl and C<sub>2</sub>-C<sub>6</sub>-alkoxycarbonyl,
- R<sup>18</sup> in each case independently of one another represent hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl or C<sub>1</sub>-C<sub>6</sub>-haloalkyl, or N(R<sup>13</sup>)(R<sup>18</sup>) represents a cycle which forms the ring M,

R<sup>19</sup> in each case independently of one another represent hydrogen or represent in each case optionally mono- or polysubstituted C<sub>1</sub>-C<sub>6</sub>-alkyl, where the substituents independently of one another are selected from the group consisting of cyano, nitro, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>2</sub>-C<sub>8</sub>-dialkylamino, CO<sub>2</sub>H, C<sub>2</sub>-C<sub>6</sub>-alkoxycarbonyl, C<sub>2</sub>-C<sub>6</sub>-alkylcarbonyl, C<sub>3</sub>-C<sub>6</sub>-trialkylsilyl and optionally substituted phenyl, where the substituents independently of one another are selected from one to three radicals W, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, phenyl or pyridyl, each of which is optionally mono- to trisubstituted by W,

M in each case represents an optionally mono- to tetrasubstituted ring which, in addition to the nitrogen atom attached to the substituent pair R<sup>13</sup> and R<sup>18</sup>, (R<sup>15</sup>)<sub>2</sub> or (R<sup>16</sup>)<sub>2</sub>, contains two to six carbon atoms and optionally additionally a further nitrogen, sulfur or oxygen atom, where the substituents independently of one another are selected from the group consisting of C<sub>1</sub>-C<sub>2</sub>-alkyl, halogen, cyano, nitro and C<sub>1</sub>-C<sub>2</sub>-alkoxy,

W in each case independently of one another represent C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>2</sub>-C<sub>4</sub>-alkenyl, C<sub>2</sub>-C<sub>4</sub>-alkynyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>2</sub>-C<sub>4</sub>-haloalkenyl, C<sub>2</sub>-C<sub>4</sub>-haloalkynyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl, halogen, cyano, nitro, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>2</sub>-C<sub>8</sub>-dialkylamino, C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino, (C<sub>1</sub>-C<sub>4</sub>-alkyl)-C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino, C<sub>2</sub>-C<sub>4</sub>-alkylcarbonyl, C<sub>2</sub>-C<sub>6</sub>-alkoxycarbonyl, CO<sub>2</sub>H, C<sub>2</sub>-C<sub>6</sub>-alkylaminocarbonyl, C<sub>3</sub>-C<sub>8</sub>-dialkylaminocarbonyl or C<sub>3</sub>-C<sub>6</sub>-trialkylsilyl,


n in each case independently of one another represent 0 or 1,

p in each case independently of one another represent 0, 1 or 2,

where, if (a) R<sup>5</sup> represents hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>2</sub>-C<sub>6</sub>-haloalkenyl, C<sub>2</sub>-C<sub>6</sub>-haloalkynyl, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio or halogen, and (b) R<sup>8</sup> represents hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>2</sub>-

C<sub>6</sub>-haloalkenyl, C<sub>2</sub>-C<sub>6</sub>-haloalkynyl, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio,  
halogen, C<sub>2</sub>-C<sub>4</sub>-alkylcarbonyl, C<sub>2</sub>-C<sub>6</sub>-alkoxycarbonyl, C<sub>2</sub>-C<sub>6</sub>-  
alkylaminocarbonyl or C<sub>3</sub>-C<sub>8</sub> dialkylaminocarbonyl, (c) then at least one  
substituent selected from the group consisting of R<sup>6</sup>, R<sup>11</sup> and R<sup>12</sup> is present  
and (d) if R<sup>12</sup> is not present, then at least one of the radicals R<sup>6</sup> and R<sup>11</sup> is  
different from C<sub>2</sub>-C<sub>6</sub>-alkylcarbonyl, C<sub>2</sub>-C<sub>6</sub>-alkoxycarbonyl, C<sub>2</sub>-C<sub>6</sub>-  
alkylaminocarbonyl or C<sub>3</sub>-C<sub>8</sub>-dialkylaminocarbonyl.

2. (Currently Amended) A The composition as claimed in according to  
claim 1 comprising at least one compound of the formula (I) in which  
W<sup>1</sup> represents hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, chlorine, bromine or  
fluorine,  
X represents C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, fluorine, chlorine or  
bromine,  
Y and Z independently of one another represent hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, halogen,  
C<sub>1</sub>-C<sub>4</sub>-alkoxy or C<sub>1</sub>-C<sub>4</sub>-haloalkyl,  
A<sup>3</sup> represents hydrogen or in each case optionally halogen-substituted C<sub>1</sub>-C<sub>6</sub>-  
alkyl or C<sub>3</sub>-C<sub>8</sub>-cycloalkyl,  
A<sup>4</sup> represents hydrogen, methyl or ethyl, or  
A<sup>3</sup> and A<sup>4</sup> furthermore together with the carbon atom to which they are attached  
represent saturated C<sub>3</sub>-C<sub>6</sub>-cycloalkyl in which optionally one ring member  
is replaced by oxygen or sulfur and which is optionally mono- or  
disubstituted by C<sub>1</sub>-C<sub>4</sub>-alkyl, trifluoromethyl or C<sub>1</sub>-C<sub>4</sub>-alkoxy,  
D represents hydrogen, in each case optionally fluorine- or chlorine-  
substituted C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>4</sub>-alkenyl or C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, or  
A<sup>3</sup> and D together represent optionally methyl-substituted C<sub>3</sub>-C<sub>4</sub>-alkanediyl in  
which optionally one methylene group is replaced by sulfur,  
G<sup>1</sup> represents hydrogen (a) or represents one of the groups



- E represents a metal ion or an ammonium ion,
- L<sup>1</sup> represents oxygen or sulfur,
- M<sup>1</sup> represents oxygen or sulfur,
- R<sup>20</sup> represents in each case optionally halogen-substituted C<sub>1</sub>-C<sub>10</sub>-alkyl, C<sub>2</sub>-C<sub>10</sub>-alkenyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, or C<sub>1</sub>-C<sub>4</sub>-alkylthio-C<sub>1</sub>-C<sub>4</sub>-alkyl or optionally fluorine-, chlorine-, C<sub>1</sub>-C<sub>4</sub>-alkyl- or C<sub>1</sub>-C<sub>2</sub>-alkoxy-substituted C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, or represents optionally fluorine-, chlorine-, bromine-, cyano-, nitro-, C<sub>1</sub>-C<sub>4</sub>-alkyl-, C<sub>1</sub>-C<sub>4</sub>-alkoxy-, trifluoromethyl- or trifluoromethoxy-substituted phenyl, or represents in each case optionally chlorine- or methyl-substituted pyridyl or thienyl,
- R<sup>21</sup> represents in each case optionally fluorine- or chlorine-substituted C<sub>1</sub>-C<sub>10</sub>-alkyl, C<sub>2</sub>-C<sub>10</sub>-alkenyl, or C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>2</sub>-C<sub>4</sub>-alkyl, represents optionally methyl- or methoxy-substituted C<sub>5</sub>-C<sub>6</sub>-cycloalkyl, or represents in each case optionally fluorine-, chlorine-, bromine-, cyano-, nitro-, C<sub>1</sub>-C<sub>4</sub>-alkyl-, C<sub>1</sub>-C<sub>4</sub>-alkoxy-, trifluoromethyl- or trifluoromethoxy-substituted phenyl or benzyl,
- R<sup>22</sup> represents optionally fluorine-substituted C<sub>1</sub>-C<sub>4</sub>-alkyl or represents optionally fluorine-, chlorine-, bromine-, C<sub>1</sub>-C<sub>4</sub>-alkyl-, C<sub>1</sub>-C<sub>4</sub>-alkoxy-, trifluoromethyl-, trifluoromethoxy-, cyano- or nitro-substituted phenyl,
- R<sup>23</sup> represents in each case optionally fluorine- or chlorine-substituted C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylamino, or C<sub>1</sub>-C<sub>4</sub>-alkylthio or represents in each case optionally fluorine-, chlorine-, bromine-, nitro-, cyano-, C<sub>1</sub>-C<sub>4</sub>-

alkoxy-, trifluoromethoxy-, C<sub>1</sub>-C<sub>4</sub>-alkylthio-, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio-, C<sub>1</sub>-C<sub>4</sub>-alkyl- or trifluoromethyl-substituted phenyl, phenoxy or phenylthio,

R<sup>24</sup> represents C<sub>1</sub>-C<sub>4</sub>-alkoxy or C<sub>1</sub>-C<sub>4</sub>-alkylthio,

R<sup>25</sup> represents C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, C<sub>3</sub>-C<sub>6</sub>-alkenyl, or C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl,

R<sup>26</sup> represents C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-alkenyl or C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, or

R<sup>25</sup> and R<sup>26</sup> furthermore together represent an optionally methyl- or ethyl-substituted C<sub>3</sub>-C<sub>6</sub>-alkylene radical in which optionally one carbon atom is replaced by oxygen or sulfur;

and ~~at least one active compound of the formula (II).~~

3. (Currently Amended) A The composition ~~as claimed in~~ according to claim 1 or 2 comprising at least one compound of the formula (I) in which

W<sup>1</sup> represents hydrogen, methyl, ethyl, chlorine, bromine or methoxy,

X represents chlorine, bromine, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy or trifluoromethyl,

Y and Z independently of one another represent hydrogen, fluorine, chlorine, bromine, methyl, ethyl, propyl, isopropyl, trifluoromethyl or methoxy,

A<sup>3</sup> represents methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, cyclopropyl, cyclopentyl or cyclohexyl,




A<sup>4</sup> represents hydrogen, methyl or ethyl, or

A<sup>3</sup> and A<sup>4</sup> furthermore together with the carbon atom to which they are attached represent saturated C<sub>6</sub>-cycloalkyl in which optionally one ring member is replaced by oxygen and which is optionally monosubstituted by methyl, ethyl, methoxy, ethoxy, propoxy or butoxy,

D represents hydrogen, represents methyl, ethyl, propyl, isopropyl, butyl, isobutyl, allyl, cyclopropyl, cyclopentyl or cyclohexyl, or

A<sup>3</sup> and D together represent optionally methyl-substituted C<sub>3</sub>-C<sub>4</sub>-alkanediyl,

G<sup>1</sup> represents hydrogen (a) or represents one of the groups



L<sup>1</sup> represents oxygen or sulfur,  
M<sup>1</sup> represents oxygen or sulfur,  
R<sup>20</sup> represents C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>2</sub>-C<sub>4</sub>-alkenyl, methoxymethyl, ethoxymethyl, ethylthiomethyl, cyclopropyl, cyclopentyl or cyclohexyl, or represents phenyl which is optionally mono- to disubstituted by fluorine, chlorine, bromine, cyano, nitro, methyl, ethyl, methoxy, trifluoromethyl or trifluoromethoxy, or represents pyridyl or thienyl, each of which is optionally mono- to disubstituted by chlorine or methyl,  
R<sup>21</sup> represents C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>2</sub>-C<sub>4</sub>-alkenyl, methoxyethyl, ethoxyethyl, or represents phenyl or benzyl,  
R<sup>25</sup> and R<sup>26</sup> independently of one another represent methyl, ethyl or together with the nitrogen atom to which they are attached represent morpholino, and ~~at least one active compound of the formula (II).~~

4. (Currently Amended) A The composition according to claim 3 as claimed in claim 1, 2 or 3 comprising at least one compound of the formula (I) in which

W<sup>1</sup> represents hydrogen or methyl,


X represents chlorine, bromine or methyl,

Y and Z independently of one another represent hydrogen, chlorine, bromine or methyl,

A<sup>3</sup> and A<sup>4</sup> furthermore together with the carbon atom to which they are attached represent saturated C<sub>6</sub>-cycloalkyl in which optionally one ring member is replaced by oxygen and is optionally monosubstituted by methyl, methoxy, ethoxy, propoxy or butoxy,

D represents hydrogen,

G<sup>1</sup> represents hydrogen (a) or represents one of the groups



L<sup>1</sup> represents oxygen or sulfur,


M<sup>1</sup> represents oxygen or sulfur,

R<sup>20</sup> represents C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>2</sub>-C<sub>4</sub>-alkenyl, methoxymethyl, ethoxymethyl, ethylmethylthio, cyclopropyl, cyclopentyl, or cyclohexyl, or represents phenyl which is optionally monosubstituted by fluorine, chlorine, bromine, methyl, methoxy, trifluoromethyl, trifluoromethoxy, cyano or nitro, or represents pyridyl or thienyl, each of which is optionally monosubstituted by chlorine or methyl,

R<sup>21</sup> represents C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>2</sub>-C<sub>4</sub>-alkenyl, methoxyethyl, ethoxyethyl, phenyl or benzyl,

R<sup>25</sup> and R<sup>26</sup> independently of one another represent methyl, ethyl or together with the nitrogen atom to which they are attached represent morpholino[[],] and at least one active compound of the formula (II).


5. (Currently Amended) A The composition according to claim 1 as claimed in claim 1, 2, 3 or 4 comprising the compound of the formula (I-a-4)



and at least one anthranilamide of the formula (II).

6. (Cancelled)

7. (Currently Amended) A The composition according to claim 1 as claimed in claim 1, 2, 3, 4, 5 or 6 comprising an anthranilamide of the formula (II-1)



in which

- R<sup>2</sup> represents hydrogen or C<sub>1</sub>-C<sub>6</sub>-alkyl,  
R<sup>3</sup> represents C<sub>1</sub>-C<sub>6</sub>-alkyl which is optionally substituted by one R<sup>6</sup>,  
R<sup>4</sup> represents C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>2</sub>-haloalkyl, C<sub>1</sub>-C<sub>2</sub>-haloalkoxy or halogen,  
R<sup>5</sup> represents hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>2</sub>-haloalkyl, C<sub>1</sub>-C<sub>2</sub>-haloalkoxy or halogen,  
R<sup>6</sup> represents -C(=E<sup>2</sup>)R<sup>19</sup>, -LC(=E<sup>2</sup>)R<sup>19</sup>, -C(=E<sup>2</sup>)LR<sup>19</sup> or -LC(=E<sup>2</sup>)LR<sup>19</sup>, where each E<sup>2</sup> independently of one another represents O, S, N-R<sup>15</sup>, N-OR<sup>15</sup>, N-N(R<sup>15</sup>)<sub>2</sub>, and each L independently of one another represents O or NR<sup>18</sup>,  
R<sup>7</sup> represents C<sub>1</sub>-C<sub>4</sub>-haloalkyl or halogen,  
R<sup>9</sup> represents C<sub>1</sub>-C<sub>2</sub>-haloalkyl, C<sub>1</sub>-C<sub>2</sub>-haloalkoxy, S(O)<sub>p</sub>-C<sub>1</sub>-C<sub>2</sub>-haloalkyl or halogen,  
R<sup>15</sup> in each case independently of one another represent hydrogen or represent in each case optionally substituted C<sub>1</sub>-C<sub>6</sub>-haloalkyl or C<sub>1</sub>-C<sub>6</sub>-alkyl, where the substituents independently of one another may be selected from the group consisting of cyano, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfinyl and C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfonyl,  
R<sup>18</sup> in each case independently of one another represents hydrogen or C<sub>1</sub>-C<sub>4</sub>-alkyl,  
R<sup>19</sup> in each case independently of one another represent hydrogen or C<sub>1</sub>-C<sub>6</sub>-alkyl,  
p independently of one another represent 0, 1, 2.

8. (Currently Amended) A The composition according to claim 1 as claimed in claim 1, 2, 3, 4, 5, 6 or 7 comprising compounds of the formula (I) (group 1) or and at least one acaricidally active compound (group 2) and at least one anthranilamide of the formula (II) in a ratio from 500:1 to 1:50.

9. (Cancelled)

10. (Currently Amended) A process for preparing a pesticide composition pesticides, comprising mixing characterized in that a synergistically effective mixture as defined in a composition according to claim 1, 2, 3, 4, 5-6 or 7 claim 1 is mixed with an extender and/or surfactant extenders and/or surfactants.
11. (Currently Amended) A method for controlling animal pests, comprising contacting an animal pest or its habitat with a composition according to claim 1 characterized in that synergistically effective mixtures as defined in claim 1, 2, 3, 4, 5-6 or 7 is allowed to act on animal pests and/or their habitat.